



US009440857B2

(12) **United States Patent**
Sorensen et al.

(10) **Patent No.:** **US 9,440,857 B2**
(45) **Date of Patent:** **Sep. 13, 2016**

(54) **PROCESS FOR HIGH-YIELD PRODUCTION OF GRAPHENE VIA DETONATION OF CARBON-CONTAINING MATERIAL**

(71) Applicant: **Kansas State University Research Foundation**, Manhattan, KS (US)

(72) Inventors: **Christopher Sorensen**, Manhattan, KS (US); **Arjun Nepal**, Manhattan, KS (US); **Gajendra Prasad Singh**, Jharkhand (IN)

(73) Assignee: **Kansas State University Research Foundation**, Manhattan, KS (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/272,924**

(22) Filed: **May 8, 2014**

(65) **Prior Publication Data**

US 2014/0335010 A1 Nov. 13, 2014

Related U.S. Application Data

(60) Provisional application No. 61/821,816, filed on May 10, 2013.

(51) **Int. Cl.**
C01B 31/04 (2006.01)

(52) **U.S. Cl.**
CPC **C01B 31/0446** (2013.01); **C01B 2204/20** (2013.01); **C01B 2204/32** (2013.01)

(58) **Field of Classification Search**
CPC C01B 31/0438; C01B 31/0446
USPC 423/445 B; 977/734
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,691,909 B2 4/2010 Sorensen et al.
2010/0278712 A1 11/2010 Swanson

FOREIGN PATENT DOCUMENTS

CN 102249219 A * 11/2011

OTHER PUBLICATIONS

Machine English translation of CN 102249219 A.*
Nepal, Arjun, Bret Flanders, and Christopher Sorensen. "Graphene in Carbon Aerosol Gels formed via Controlled Hydrocarbon Detonation." (2011).
Sorensen, Christopher M., et al. "Formation of Light-weight Low-density Materials via Gas Phase Aerosol Gelation." MRS Proceedings. vol. 1306. Cambridge University Press, 2011.*
The Power Point presentation entitled "Formation of Light-weight Low-density Materials via Gas Phase Aerosol Gelation," presented at AAAR 30th Annual Conference, 2011.

* cited by examiner

Primary Examiner — Richard M Rump

(74) *Attorney, Agent, or Firm* — Hovey Williams LLP

(57) **ABSTRACT**

A method of producing pristine graphene particles through a one-step, gas-phase, catalyst-free detonation of a mixture of one or more carbon-containing compounds hydrocarbon compounds and one or more oxidizing agents is provided. The detonation reaction occurs very quickly and at relatively high temperature, greater than 3000 K, to generate graphene nanosheets that can be recovered from the reaction vessel, such as in the form of an aerosol. The graphene nanosheets may be stacked in single, double, or triple layers, for example, and may have an average particle size of between about 35 to about 250 nm.

31 Claims, 10 Drawing Sheets

